

Silicone Therapy

Silicone treatment is a widely used therapy acting as an agent to soften flatten and pale active scars. There are many opinions on how it works but none are conclusive at this time. There are many suggestions that it creates a barrier to water loss (this barrier is damaged during the injury to the skin) which prevents the scar from becoming dry and ultimate aids the maturation process.

There are many silicone products on the market in a variety of forms all aiming to achieve the same outcome:

- Cream / gel
- Sheets
- Spray

The application of each of these gels has specific wearing instructions so see the manufacturer's information sheets for details.

Sheet Silicones: The sheet silicones last between 7 days and 6 weeks. You should cut the sheet to the size of your scar, leaving at least a centimetre border around the edge and re-apply the same piece until it is no longer adhesive. You can normally wash the sheet with soapy water and leave it to air dry (so have two pieces in use to allow the silicone to fully dry before repeat application).

When using sheet silicone you need to slowly build up the wearing time (you can wear it full time, just removing to apply any moisturiser or to cleanse the area) so for you first week we recommend a 4 hour build up time:

Day 1	Day 2	Day 3	Day 4	Day 5	Day 6
4 hours	8 hours	12 hours	16 hours	20 hours	Up to 24 hours

Silicone Creams / gels and Sprays: These are normally applied and massaged into the scar twice daily. Apply your moisturiser first, when this is fully absorbed then apply you silicone cream sparingly. If you are applying to an area that is washed regularly e.g. your hand then you may need to repeat more than twice a day. Apply a test patch first and leave 24 hours before you start using to check your skin does not react to the silicone. Make sure you fully cleanse your skin between each application.

Allergies Precautions: It is possible to be allergic to silicone gels/sheets so always test your silicone first. If you are allergic then the area may become red, itchy and may develop a rash. If this happens, then stop using the silicone and allow the skin to settle. Please contact your therapist to discuss. Another cause of a reaction can be that the skin and your gel (if using a sheet) are not thoroughly cleansed between applications.

It is also advised that you **do not** use silicone on open wounds.

Treatment Regime	after moisturiser			,	Sheets: can be worn full time, removed to moisturise and cleanse the scar				
Products	A variety of products are available – see some examples below								
Precautions	Gels: Always test and wait 24 hours before using fully Sheets: Use suggested build up time If your scar becomes more itchy, painful or develops a rash then remove and contact your therapist								
Product examples:	Gel	Sheet	Spray	Size		Where to purchase / Prices 2014			
Scarsil	√			15m	l or 30ml	www.jobskin.co.uk 0115 9734300 £25 -£35.00			
Scarfix		√		25.5 3.75 Brea 7.5c Brea	n x 20cm ccm x 30.5cm ccm x 22.5cm est Circle: m diameter est Piece: ccm x 14.5cm	www.jobskin.co.uk 0115 9734300 £16.00 - £31.00			
Dermatix fabric		√		13 x 13 x	13cm 13cm 25cm 30cm	www.dermatix.co.uk 0800 7834995 £13.09 - £98.94			
Mepiform		√		_	7cm 18cm 31cm				
Silderm	√		√		6g or 15g ay 30ml or 60ml	www.silderm.com 01260 271666 £9.99 to £52.00			